CORRECTION

Open Access



Correction to: Role of hepatobiliary scintigraphy with different semi-quantitative parameters along with histopathological scoring in differentiating biliary atresia from neonatal hepatitis

Nsreen R. A. Mohamadien¹, Rania Makboul², Shereen M. Galal³ and Nadia M. Mostafa^{1*}

Correction to: Egyptian Journal of Radiology and Nuclear Medicine (2021) 52:111 https://doi.org/10.1186/s43055-021-00482-5

Following the publication of the original article [1], the authors identified that the captions were incorrectly assigned to the figures and the word **cholestasis** consisted of a space.

The changes have been highlighted in **bold typeface**.

The original article [1] has been corrected.

Fig. 1 Liver core stained by H&E of a case of BA showing periportal fibrosis (×40). B &C: Higher power view of previous case showing bile ductular proliferation, bile plugs, portal fibrosis and neutrophilic infiltration (×200,×400 respectively) D: Liver core stained by H&E of case of NH showing periportal and lobular inflammation (×100). E: higher power of previous case showing lobular inflammation and lobular disarray (×200). F: High power of a case of NH showed cholate stasis, lobular inflammation and giant cell formation (×400)

Fig. 2 Association between histopathological scoring and diagnosis of BA

The original article can be found online at https://doi.org/10.1186/s43055-021-00482-5.

¹ Department of Clinical Oncology and Nuclear Medicine, Faculty of Medicine, Assiut University, Assiut, Egypt

Full list of author information is available at the end of the article

Den Springer Open

© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Fig. 6 Association between semi-quantitative parameters and BA

Author details

¹Department of Clinical Oncology and Nuclear Medicine, Faculty of Medicine, Assiut University, Assiut, Egypt. ²Department of Pathology, Faculty of Medicine, Assiut University, Assiut, Egypt. ³Department of Pediatrics, Faculty of Medicine, Assiut University, Assiut, Egypt.

Published online: 24 August 2021

Reference

 Mohamadien NR et al. Role of hepatobiliary scintigraphy with different semi-quantitative parameters along with histopathological scoring in differentiating biliary atresia from neonatal hepatitis. Egypt J Radiol Nuclear Med.2021;52:111. https://doi.org/10.1186/s43055-021-00482-5.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

^{*}Correspondence: nadia.khalifa@med.aun.edu.eg; drnadia1980@gmail.com